

Casimir effect around disclinations

Fernando Moraes*
School of Natural Sciences
Institute for Advanced Study
Princeton, NJ 08540
U. S. A.

Abstract

This communication concerns the structure of the electromagnetic quantum vacuum in a disclinated insulator. It is shown that a nonzero vacuum energy density appears when the rotational symmetry of a continuous insulating elastic medium is broken by a disclination. An explicit expression is given for this Casimir energy density in terms of the parameter describing the disclination.

*On leave from:
Departamento de Física
Universidade Federal de Pernambuco
50670-901 Recife, PE, Brazil

1 Introduction

Space-time being isotropic and homogeneous implies that the vacuum state of a quantum field must be invariant under translations or rotations. The presence of external fields, a change in the geometry (curvature, torsion) or a change in the topology (defects, boundaries) of the space breaks the underlying symmetry of space-time, inducing changes in the vacuum state of the quantum field as compared to its field-free, unbound, Minkovski space counterpart. This is the celebrated Casimir effect [1, 2].

Topological defects certainly offer an interesting (and quite unexplored) arena for the study of vacuum polarization phenomena in condensed matter systems. In previous publications [3, 4] we studied the non-relativistic quantum mechanics of electrons and holes in the presence of a disclination in an insulating elastic medium, with and without an external magnetic field. In the present work, I study the Casimir effect of the electromagnetic field in an infinite medium of constant dielectricity ε and permeability $\mu = 1$ (a nonmagnetic medium), with a disclination at the origin of the coordinate system, at zero temperature. It is investigated the electromagnetic Casimir effect since it is more likely to be experimentally observable in the solid state than the fermionic case [5]. This is due to the fact that massive fields, like the fermion field, give Casimir energies exponentially decreasing with their mass. An estimate [5] for electrons confined to regions of nanometric s

ize, i.e. $\sim 10^{-7}$ cm, gives an exponential damping factor for the Casimir energy of $\sim 5 \times 10^3$. While the electromagnetic Casimir energy for light confined to a similar region is $\sim 10\text{eV}$, the electronic effect is then roughly $e^{-5 \times 10^3}$ weaker. Notice that while reference [5] deals with the Casimir effect due to confinement to mesoscopic regions, in this work the interest is in the effect outside a defect - no confinement involved. In any case, one should expect a much weaker electronic Casimir effect, then justifying the choice of presenting the electromagnetic effect here.

2 QED in conical space

Quantum field theory has been extensively studied in the conical space-time of cosmic strings [6]. The large body of information available on this subject makes an excellent guide for the study of quantum field theoretical aspects of disclinations in solids, in the context of the theory of defects/three-dimensional gravity of Katanaev and Volovich [7].

Disclinations, like cosmic strings, are associated with a conical metric [7] whose spatial part is given, in cylindrical coordinates, by

$$ds^2 = dz^2 + d\rho^2 + \alpha^2 \rho^2 d\varphi^2. \quad (1)$$

With $\alpha = 1 + \frac{\lambda}{2\pi}$, the disclination is obtained by either removing ($\lambda < 0$) or inserting ($\lambda > 0$) a wedge of material, of dihedral angle λ , to the otherwise smooth elastic continuum. The resulting space, described by the above metric, has a null curvature tensor everywhere, except at the defect where it has a two-dimensional δ -function singularity given by [8]

$$R_{12}^{12} = R_1^1 = R_2^2 = 2\pi \frac{1-\alpha}{\alpha} \delta_2(\rho), \quad (2)$$

where $\delta_2(\rho)$ is the two-dimensional delta function in flat space. It is clear that $0 \leq \alpha < 1$ corresponds to a positive-curvature disclination and $\alpha > 1$ to a negative-curvature disclination. $\alpha = 1$, of course, describes the Euclidean medium (absence of disclinations).

The relevant physical quantity in the study of the Casimir effect is the vacuum expectation value of the energy-momentum tensor, or stress tensor, $\langle \hat{T}_\mu{}^\nu \rangle$. Since this tensor diverges, in order to obtain physically meaningful results, one should subtract from it its also divergent flat space version¹. That is, one should calculate the renormalized vacuum expectation value

$$\langle \hat{T}_\mu{}^\nu \rangle_{ren} = \langle \hat{T}_\mu{}^\nu \rangle_{discl} - \langle \hat{T}_\mu{}^\nu \rangle_{flat}. \quad (3)$$

¹In many cases an additional regularization is needed to ensure a finite result.

In this work, I will be concerned only with the $\langle \hat{T}_0^0 \rangle$ component, since it is the one that gives the energy density.

Being the electromagnetic field a gauge field, care must be exercised in order that appropriate gauge-fixing be taken into account in the quantum theory. As it turns out [9] the convenient gauge here is

$$\nabla_i A^i = \nabla_a A^a = 0 \quad (i = 0, 1; a = 2, 3), \quad (4)$$

where A^μ is the electromagnetic vector potential. In this gauge the energy density assumes the very simple form

$$\langle \hat{T}_0^0 \rangle = i\hbar c \lim_{x' \rightarrow x} \nabla_i \nabla^i G(x - x'), \quad (5)$$

where $G(x - x')$ is the scalar Green's function, which in the absence of the defect is

$$G_0(x - x') = -\frac{i}{4\pi^2 \varepsilon} \frac{1}{(x - x')^2}. \quad (6)$$

$x - x'$ is the space-time interval between x and x' , its square being given by

$$(x - x')^2 = -c^2(t - t')^2 + (z - z')^2 + \rho^2 + \rho'^2 - 2\rho\rho' \cos(\varphi - \varphi'). \quad (7)$$

The energy density is then

$$\langle \hat{T}_0^0 \rangle_{ren} = i\hbar c \left(-\frac{1}{c^2} \partial_t^2 + \partial_z^2 \right) (G_\alpha(x - x') - G_0(x - x')) \big|_{x'=x}, \quad (8)$$

where $G_\alpha(x - x')$ is the scalar Green's function in the medium with boundary conditions. Due to the disclination, the boundary condition in this problem is a periodicity of $2\pi\alpha$ in the angular variable φ . Accordingly, imposition of this condition on the free space Green's function can be done by reperiodising it via the contour integral [10]

$$G_\alpha(x - x') = \frac{-i}{4\pi\alpha} \int_A d\gamma G_0(\varphi - \varphi' - \gamma) \frac{\exp(i\gamma/2\alpha)}{\sin(\gamma/2\alpha)}. \quad (9)$$

The singularities in the integrand are poles that can be easily found by writing

$$(x - x')^2 = 2\rho\rho'[\cosh \beta - \cos(\varphi - \varphi')],$$

where

$$\cosh \beta = \frac{-c^2(t - t')^2 + (z - z')^2 + \rho^2 + \rho'^2}{2\rho\rho'}.$$

Since $\cosh \beta = \cos i\beta$, the poles of $G_0(\varphi - \varphi' - \gamma)$ are at $\gamma = \pm i\beta + \varphi - \varphi' + n2\pi$. On the other hand, $\sin(\gamma/2\alpha)$ contributes with poles at $\gamma = n2\pi\alpha$. In both cases $n \in \mathbb{Z}$.

As shown in figure 1, the contour A has two branches, one in the upper half γ -plane from $(\pi + \varphi - \varphi') + i\infty$ to $(-\pi + \varphi - \varphi') + i\infty$ passing below the singularity $\gamma = i\beta + \varphi - \varphi'$ and the other in the lower half-plane from $(-\pi + \varphi - \varphi') - i\infty$ to $(\pi + \varphi - \varphi') - i\infty$ passing above the singularity at $\gamma = -i\beta + \varphi - \varphi'$.

In the limit $x' = x$ the poles at $\gamma = \pm i\beta + \varphi - \varphi' + n2\pi$ will be lying at $\gamma = n2\pi$. So, one is left with poles at $\gamma = n2\pi$ and $\gamma = n2\pi\alpha$. Considering this, there is a deformation [10] of the contour A that will prove to be quite convenient. Just deform it into an anticlockwise loop B around the pole at $\gamma = 0$ and the two vertical lines, Γ , $\gamma = b + iy$ and $\gamma = -b + iy$, with $-\infty < y < \infty$ and b a constant (see figures 2 and 3). This can only be done if the poles lying on the $\text{Re}\gamma$ -axis, other than the one at $\gamma = 0$, are outside the region limited by the lines; i.e., b must be chosen to be less than the smaller of 2π and $2\pi\alpha$. As shown in the appendix, the contribution from loop B turns out to be exactly $G_0(x - x')$. Since renormalization requires $G_\alpha(x - x') - G_0(x - x')$, all one has to consider is the contribution from the two vertical lines Γ . Equation (8) becomes then

$$\langle \hat{T}_0^0 \rangle_{ren} = \frac{-i\hbar c}{4\pi\alpha} \int_{\Gamma} d\gamma \frac{\exp(i\gamma/2\alpha)}{\sin(\gamma/2\alpha)} \left(-\frac{1}{c^2} \partial_t^2 + \partial_z^2 \right) G_0(\varphi - \varphi' - \gamma) \big|_{x'=x}. \quad (10)$$

It is easily found that

$$\left(-\frac{1}{c^2} \partial_t^2 + \partial_z^2 \right) G_0(\varphi - \varphi' - \gamma) \big|_{x'=x} = \frac{i}{16\pi^2 \varepsilon \rho^4 \sin^4(\gamma/2)}. \quad (11)$$

Furthermore, using the symmetry of the contour Γ to replace the exponential by a cosine in equation (10) and then deforming the contour into a clockwise loop around the origin leads to

$$\langle \hat{T}_0^0 \rangle = \frac{i\hbar c}{64\pi^3 \varepsilon \rho^4 \alpha} \oint d\gamma \frac{\cot(\gamma/2\alpha)}{\sin^4(\gamma/2)}. \quad (12)$$

This integral can be easily evaluated using calculus of residues, by expanding the integrand in powers of γ such that the term of order γ^{-1} gives $\oint \frac{d\gamma}{\gamma} = -2\pi i$. The expansion results in

$$\frac{\cot(\gamma/2\alpha)}{\sin^4(\gamma/2)} = 32\alpha\gamma^{-5} + \left(\frac{16}{3}\alpha - \frac{8}{3\alpha}\right)\gamma^{-3} + \left(\frac{22}{45}\alpha - \frac{4}{9\alpha} - \frac{2}{45\alpha^3}\right)\gamma^{-1} + O(\gamma^0). \quad (13)$$

The energy density is then

$$\langle \hat{T}_0^0 \rangle = \frac{\hbar c}{720\pi^2 \varepsilon \rho^4} \left(11 - \frac{10}{\alpha^2} - \frac{1}{\alpha^4}\right). \quad (14)$$

This result has been found by a variety of ways in the context of cosmic strings [9, 11, 12] and also for a metallic wedge [13].

Notice that in the flat space limit, $\alpha = 1$, $\langle \hat{T}_0^0 \rangle_{ren} = 0$ as it should. Also, although values of $\alpha > 1$ are probably not physically sensible for cosmic strings because it would correspond to negative linear mass densities, in condensed matter it describes negative-curvature disclinations [3, 4]. The immediate consequence for condensed matter systems with such defects is a dependence of the sign of the force exerted on the defect by the electromagnetic vacuum, on the sign of the defect curvature. This is certainly important for the study of the stability of the defects. The obvious implication is that a disclination dipole, or edge dislocation [7], is more stable than either disclination standing by itself and that the interaction between the pair is attractive. This is a result of the vacuum polarization of the electromagnetic field only. The total interaction and overall stability depend on contributions from all the relevant fields.

3 Concluding remarks

The purpose of this Letter is to draw attention to the importance of quantum field theoretical effects in condensed matter enclosing topological defects. A very simple case has been considered: the medium is infinite (e.g. no external boundaries were considered), nonmagnetic, has only one defect (disclination), its dielectricity is frequency-independent and the temperature set to $T = 0$. Of course, these oversimplifications give a distorted picture of the real phenomenon. Nevertheless, it is shown that a finite electromagnetic Casimir energy density is present in the outer region of the disclination. It would be interesting to extend the present results to finite temperatures and to magnetic media. Moreover, use of a frequency-dependent complex dielectric constant would not only provide a more realistic calculation for the Casimir energy but also one that takes into account absorption of energy inside the dielectric, something that happens in real materials.

Acknowledgements

This work was partially supported by CNPq.

APPENDIX

The anticlockwise loop integral

$$\frac{-i}{4\pi\alpha} \oint_B d\gamma \frac{\exp(i\gamma/2\alpha)}{\sin(\gamma/2\alpha)} G_0(\varphi - \varphi' - \gamma)$$

can be evaluated in the same way as the integral in equation (12). Again, symmetry of the contour permits the substitution of the exponential by a cosine. The resulting $\cot(\gamma/2\alpha)$ expanded around $\gamma = 0$ is

$$\frac{2\alpha}{\gamma} - \frac{\gamma}{6\alpha} - \frac{\gamma^3}{360\alpha^3} + O(\gamma^5),$$

implying

$$\frac{-i}{4\pi\alpha} \oint_B d\gamma \frac{\exp(i\gamma/2\alpha)}{\sin(\gamma/2\alpha)} G_0(\varphi - \varphi' - \gamma) = \frac{-i}{2\pi} \oint_B \frac{d\gamma}{\gamma} G_0(\varphi - \varphi' - \gamma) = G_0(x - x').$$

References

- [1] V.M. Mostepanenko and N.N. Trunov, Sov. Phys. Usp. 31 (1988) 965.
- [2] G. Plunien, B. Müller and W. Greiner, Phys. Rep. 134 (1986) 87.
- [3] Claudio Furtado and Fernando Moraes, Phys. Lett. A 188 (1994) 394.
- [4] C. Furtado, B.G.C. da Cunha, F. Moraes, E.R. Bezerra de Mello and V.B. Bezerra, Phys. Lett. A 195 (1994) 90.
- [5] I.H. Dur and M. Tomak, Phys. Lett. A 176 (1993) 265.
- [6] See for example, G.W. Gibbons, S.W. Hawking and T. Vachaspati, eds., The formation and evolution of cosmic strings (Cambridge Univ. Press, Cambridge, 1990), part 4 and references therein.
- [7] M.O. Katanaev and I.V. Volovich, Ann. Phys. (N.Y.) 216 (1992) 1.
- [8] D.D. Sokolov and A.A. Starobinskii, Sov. Phys. Dokl. 22 (1977) 312.
- [9] V.P. Frolov and E.M. Serebriany, Phys. Rev. D 35 (1987) 3779.
- [10] J.S. Dowker, Phys. Rev. D 36 (1987) 3095.
- [11] M.R. Brown, A.C. Ottewill and D.N. Page, Phys. Rev. D 33 (1986) 2840.
- [12] J.S. Dowker, Phys. Rev. D 36 (1987) 3742.
- [13] D. Deutsch and P. Candelas, Phys. Rev. D 20 (1979) 3063.

Figure Captions

Figure 1: Integration contour for equation (9).

Figure 2: Intermediate deformation of the contour.

Figure 3: Final form of the deformed contour.




Figure 1




Figure 2




Figure 3